INDEX TO VOLUME 41

New names and the final members of new combinations are in **bold face type.**

Abies, 625; alba, 628; americana, 627, 628, 629; balsamea, 627, 628; canadensis, 627, 628, 629; mariana, 627, 628; nigra, 628
Abrus precatorius, 107
Abutilon Lemmoni, 303
Acacia Berteriana, 8; lentiscifolia, 8; suffrutescens, 303
Acaena, 323, 324; agrimonioides, 323, 324; californica, 324, 325; elongata, 323, 324; elongata gracilis, 324; elongata robusta, 324; laevigata, 323, 324; lappacea, 323, 324; pinnatifida, 324; torilicarpa, 324; tridactyla, 324; trifida, 324
Acalypha alopecuroidea, 12
Acanthospermum hispidum, 13
Acer floridanum, 214, 219; rubrum, 559; saccharum, 512
Acetabula, 158
Achillea borealis, 102; lanulosa, 100; Millefolium, 355, 358, 512, 514, 516, 517, 519, 520; multiflora, 102; subalpina, 100, 471
Achroanthes unifolia, 29, 32, 35
Acolea crenulata, 612
Acomastylis, 490, 491, 494; depressa, 494; gracilipes, 494, 495; humilis, 494; Rossii, 100, 494; sericea, 100, 469, 471, 494, 495; turbinata, 100, 463, 469, 470, 495
Adenostoma, 330, 331; brevifolium, 330; fasciculatum, 330; fasciculatum densifolium, 330; fasciculatum hirsutum, 330; fasciculatum obtusifolium, 330; sparsifolium, 331
Adiantum Capillus-veneris, 44, 308; **rimicola,** 308
Aesculus octandra, 29, 34, 38; Pavia, 214
Aethusa Cynapium, 84
Afzelia cassioides, 561; macrophylla, 29, 35, 45; pectinata, 212
Agaricus, 179; campestris, 111
Agrimonia, 325; Bicknellii, 328; Brittoniana, 329; Eupatoria, 325, 326, 328; Eupatoria glabra, 326; Eupatoria hirsuta, 326; Eupatoria mollis, 327; Eupatoria parviflora, 329; gryposepala, 326; incisa, 329, 330; macrocarpa, 326; microcarpa, 327, 329; mollis, 327; parviflora, 326, 329, 330; platycarpa, 327, 329; polyphylla, 329; Pringlei, 327, 329; pubescens, 327, 328; pumila, 327; rostellata, 326, 328, 329; serrifolia, 329; striata, 325, 327, 328, 329; suaveolens, 329
Agropyron andinum, 99, 463, 464; biflorum, 96, 99, 469; caninum, 96, 469; latiglume, 99, 463, 464; repens, 514, 516, 518; Scribneri, 101, 463
Agrostis, 359, 362; alba, 359, 360, 512, 514, 515, 516, 518; hyemalis, 470; perennans, 308
Akebia quinata, 367
Alaska, Observations on the edge of the forest in the Kodiak region of, 381; Report on the Hepaticae of, 577
Alchemilla, 319, 320; achilleaefolia, 322; alpina, 319; domingensis, 321; filicaulis, 320; glomerulans, 320; hirsuta campestris, 321; nivalis, 321; orbiculata, 320; pectinata, 320; pinnata, 321, 322; pratensis, 320; sibbaldiaefolia, 321; speciosa, 222; vulgaris, 320; Wichurae, 320
Algites americana, 297
ALLARD, H. A., A review of investigations of the mosaic disease of tobacco, together with a bibliography of the more important contributions, 435
Allium, 358; canadense, 358; cernuum, 356, 358
Allocarya hispidula, 341
Alnus rubra, 580; rugosa, 560; sinuata, 472, 473; tenuifolia, 338, 340
Aloe ferox, 126
Alopecurus aristulatus, 470, 471
Alsine americana, 99, 465, 466; baicalensis, 467; dichotoma, 99; laeta, 97, 99, 467, 470; longipes, 97; polygonoides, 99, 468, 469
Alsinopsis obtusiloba, 462, 463; propinqua, 95, 467; quadrivalvis, 96, 468, 469; Rossii, 96, 468, 469; verna, 95, 96
Althea officinalis, 127
Amaranthus, New or notable species of, 505
Amaranthus, 505, 507, 508, 509; acutilobus, 509; blitoides, 512, 517; bracteosus, 508; crassipes, 509; emarginatus, 509; graecizans, 512; Greggii, 507; hybridus, 509; **myrianthus,** 506;

Palmeri, 506; pubescens, 508; retroflexus, 505, 510, 514, 516; Torreyi, 507; Torreyi suffruticosus, 505, 506; viscidulus, 508, 509; **Watsoni,** 505
Amarella Amarella, 96; arctophila, 102; monantha, 96; plebeja, 96, 470; propinqua, 102; scopulina, 472; strictiflora, 470, 472; tenella, 96
Ambrosia artemisiifolia, 357, 360, 512, 514; psilostachya, 512
Amelanchier Cusickii, 341; florida, 340; sp. 214
Amomum Melegueta, 107, 127
Amorpha canescens, 355, 356
Ananas sativa, 153
Anastrepta, 588, 589; orcadensis, 588, 589, 601, 610, 614, 615
Anastrophyllum Reichardtii, 613, 615
Anastrophus compressus, 560
Anchistea virginica, 560
Ancistrum, 323
Andromeda Cooki, 298; grandifolia, 298; novae-caesareae, 298; Parlatorii, 298
Andropogon furcatus, 354, 355, 357, 363; scoparius, 354, 357; virginicus, 28, 33, 35
Androsace albertina, 96; carinata, 96, 462, 464; Chamaejasme, 96; septentrionalis, 102; subumbellata, 470
Anemiopsis, 232
Anemone, 490; canadensis, 29, 35; globosa, 101; narcissiflora, 96; parviflora, 94; tetonensis, 101; zephyra, 96, 470
Anethum graveolens, 84
Aneura major, 612
Angelica atropurpurea, 107; Grayi, 101, 470; Roseana, 471
Anisacanthus Thurberi, 303
Anneslia **colletioides,** 18; **enervis,** 18
Antennaria alpina, 97, 102; aprica, 463; media, 97, 463; monocephala, 102; nardina, 472; plantaginifolia, 356, 357; rosea, 339; sedoides, 465; umbrinella, 97
Anthelia julacea, 601; Juratzkana, 613
Anthemis Cotula, 514, 516; nobilis, 126
Anthonomus grandis Thurberiae, 301
Anthropogon barbellatum, 99, 468; ciliatum, 99
Anticlea coloradensis, 466
Antiphylla oppositifolia, 466
Antrophyum minimum, 398; Wercklеanum, 398
Aphanes, 319, 320, 321; arvensis, 320; australis, 320; cuneifolia, 320; macrosepala, 320; occidentalis, 320
Aphyllon, 231
Apium graveolens, 107, 127; Petroselinum, 107, 127
Apocynum androsaemifolium, 109, 340; cannabinum, 109, 359
Appearance of polar bodies in the spermogenous tissue of Ricciocarpus natans (L.) Corda, The, 333
Aquilegia brevistyla, 100; coerulea, 467; flavescens, 470; Jonesii, 464; saximontana, 100, 465, 466
Aragallus deflexus, 100, 470; foliolosus, 100; Hallii, 464; podocarpus, 97, 464
Aralia groenlandica, 298; nudicaulis, 83; racemosa, 222; Ravniana, 298; spinosa, 28, 32, 35, 41, 42, 44, 46, 214, 560; Washingtoniana, 296
Araliaceae, 83
Araliopsis, 296, 297; breviloba, 296; cretacea, 296; cretacea dentata, 296; cretacea salisburiaefolia, 297
Araucaria bladenensis, 297; marylandica, 297
Archangelica officinalis, 126
Arctium Lappa, 121; minus, 514
Arctostaphylos Uva-ursi, 415
Arctuos, 472
Arenaria Fendleri, 464; serpyllifolia, 514, 515, 520; Tweedyi, 464
Aristida dichotoma, 29, 36; stricta, 561, 563
Aristolochia reticulata, 107; Serpentaria, 107, 127
Arnica cordifolia, 339; montana, 107, 126; Parryi, 471
Aronia nigra, 29, 35, 421
Artemisia, 110; arctica, 102; borealis, 96; ludoviciana, 338, 340, 512; norvegica, 96; Parryi, 101, 468; Pattersonii, 101, 468; pauciflora, 107; Richardsoniana, 102; saxatilis, 468; saxicola, 96; scopulorum, 101, 467; spithamaea, 96, 468, 469
Aruncus Aruncus, 28, 33, 35
Asarum arifolium, 214, 216
Asclepias, 109; incarnata, 358; linearis, 303; mexicana, 340; Sullivantii, 29, 32, 36, 38, 356, 358; syriaca, 358; tuberosa, 109, 355, 356; variegata, 29, 32, 34, 38; verticillata, 355, 360
Ascobolus, 161, 172, 179, 183, 195, 196; carbonarius, 159, 160, 162, 163, 164, 166, 172, 187, 192, 193, 194, 197; furfuraceus, 159, 160, 161, 179, 180, 181, 183, 184, 186, 187, 188, 189, 193, 194, 195; glaber, 160, 193; immersus, 186, 190, 194; Leveillei, 189, 190, 194; magnificus, 181, 183, 185, 186, 192, 193, 196, 197; mirabilis, 162, 166; pusillus, 193; Winteri, 186, 189, 194
Ascodesmis, 161, 181, 190, 192, 193; nigricans, 159, 184
Ascophanus carneus, 159, 160, 172, 193; pilosus, 159
Ascyron hypericoides, 28; multicaule, 33, 35

Asimina parviflora, 212
Aspergillus, 169, 171, 182, 193; flavus, 289; herbariorum, 159, 161; niger, 290; repens, 159
Asperula, 225
Aspidiophyllum, 296, 297; trilobatum, 297
Asplenium cecilensis, 297; Dicksonianum, 296; montanum, 29, 32, 36; pinnatifidum, 28, 33, 36
Aster, 13; adnatus, 13, 14; alpinus, 94, 471; apricus, 470; **bahamensis,** 13, 14; Bracei, 13 15; **Burgessii,** 13, 14; carneus, 15; divaricatus, 28, 33, 35; Douglasi, 338; dumosus, 13, 15; expansus, 15; exilis, 13, 15; foliaceus frondosus, 338; **Grisebachii,** 13, 14; inconspicuus, 13, 15; laevis Geyeri, 338, 340; lucayanus, 13, 14; macrophyllus, 29, 35; marginatus, 14; multiflorus, 354, 360; Novae-Angliae, 357; oreganus, 339, 340; panniculatus, 359; sericeus, 354; squamatus, 13, 15; Trifolium, 15; tripolium, 15
Asterella fragrans, 613; tenella, 613
Astragalus alpinus, 93; Purshii, 340; villosus, 212
Atelophragma elegans, 470
Atlantic coastal plain, Contributions to the Mesozoic flora of the, X—Maryland, 295
Atriplex angustifolia, 303
Atropa Belladonna, 107, 125
ATWELL, R. S. The appearance of polar bodies in the spermogenous tissue of Ricciocarpus natans (L.) Corda, 333
Avena, 225; americana, 100; Mortoniana, 100, 464; sativa, 516, 520
Avicennia, 231
Azalea glauca, 413; hispida, 413; lutea, 29, 30, 32, 35; nitida, 413; viscosa, 413

Bacillus mycoides, 290; prodigiosus, 290; pyocyaneus, 290; subtilis, 290; Tetani, 267
Baeomyces roseus, 159
BAILEY, V. Wild cotton plant (Thurberia thespesioides) in Arizona, The, 301
Baldwinia uniflora, 561
Balsamia, 182
Balsamorhiza sagittata, 342
Baptisia bracteata, 355; perfoliata, 563
Barosma betulina, 107, 110
Batodendron arboreum, 212, 214
Batrachospermum, 161, 167, 171, 172, 173
Bauhinia marylandica, 297
Bazzania deflexa, 598; Didericiana, 597; Pearsoni, 596, 597, 615, 616; tricrenata, 597, 598; trilobata, 598
Belairia **parvifoliola,** 19
BENEDICT, R. C. Revision of the genus Vittaria, J. E. Smith, A, 391
Berberis Aquifolium, 127; repens, 341
Bernhardia dichotoma, 56; ramulosa, 57
BERRY, E. W. Contributions to the Mesozoic flora of the Atlantic coastal plain, X.—Maryland, 295
Besseya alpina, 100, 470; plantaginea, 464; plantaginifolia, 100
Betula glandulosa, 472, 473; lutea, 28, 32, 35, 44; nigra, 28, 33, 35; papyrifera alaskana, 381
BICKNELL, E. P. The ferns and flowering plants of Nantucket,—XII, 71; XIII, 411
Bidens, aristosa, 30, 35
Bignonia crucigera, 214, 560
Bistorta bistortoides, 100, 470, 472; linearifolia, 100, 470; vivipara, 93, 472
BLACK, C. A. Branched cells in the prothallium of Onoclea sinsibilis L., 617
Black-soil prairie station in northeastern Illinois, A., 351
Blasia pusilla, 612
Blephariglottis lacera, 29, 34, 38; peramoena, 29, 31, 35
Blepharoneuron tricholepis, 470
Blepharostoma setiforme, 613; trichophyllum, 601
Boletus, 179
Bonnemaisonia, 167
Bonnetia, 20; anceps, 20
Botrytis cinerea, 290; vulgaris, 290
Bourreria, 10; grandiflora, 11; **moaensis,** 10; **mucronata,** 10; **Nashii,** 11; pauciflora, 11; setoso-hispida, 10
BOWMAN, H. H. M. Mechanical tissue development in certain North American vines, 365
Brachyphyllum macrocarpum, 297; macrocarpum formosum, 297
Brachysiphon, 223
Branched cells in the prothallium of Onoclea sensibilis L., 617
Brassica arvensis, 514, 517
Brauneria pallida, 355; purpurea, 30
Brickellia californica, 303
BRITTON, N. L. Studies of West Indian plants—V, 1
Bromus Kalmii, 356; secalinus, 514
Bryonia alba, 126; dioica, 126
Bryopteris, 608
Bumelia lanuginosa, 214; praenuntia, 298
Bupleurum americanum, 100; purpureum, 100; ranunculoides, 100
Burmannia, 229; candida, 230; Championii, 222, 230; coelestis, 230
Burshia, 497

Cactaceae, 71
Cactus missouriensis, 47; Opuntia, 71

Calamagrostis arundinacea, 95; canadensis, 97, 358, 359; Langsdorfii, 97, 471; purpurascens, 95, 463
Calliandra colletioides, 18; erophylla, 303; prehensilis, 8; revoluta, 5; Saman, 8
Callicarpa americana, 214, 560; Hitchcockii, 12
Callistemon, 153
Callithamnion, 174, 177
Callitriche palustris, 474
Calluna vulgaris, 415, 416, 417; vulgaris pubescens, 417
Calosiphonia, 176
Caltha biflora, 99; cheledoni, 99, 472, 473; leptosepala, 99, 472, 473; rotundifolia, 99, 472, 473
Calypogeia Neesiana, 596; Trichomanis, 596
Campanula americana, 363; lasiocarpa, 102; Parryi, 470; petiolata, 470; rotundi rotundifolia, 339; uniflora, 93, 94, 470
Camptosorus rhizophyllus, 617
Campulosus aromaticus, 560, 565
Cannabis sativa, 107, 121
Capnoides sempervirens, 29, 35
Capsella Bursa-pastoris, 514, 515, 516, 517, 518
Capsicum fastigiatum, 107
Cardamine bellidifolia, 102; rotundifolia, 30
Carduus undulatus, 341; virginicus, 30
Carex, 471; Clarkii, 298; costellata, 28, 35; floridana, 214; sp., 357
Carphephorus Pseudo-Liatris, 561, 563, 567
Carpinus caroliniana, 559
Carpites liriophylli, 297
Carpolithus septloculus, 298
Carthamus tinctorius, 126
Carum Carvi, 107, 127
Cassandra, 421
Cassia acutifolia, 107; angustifolia, 107; Chamaecrista, 354, 360; nictitans, 28, 35
Cassiope Mertensiana, 98, 467; tetragona, 98, 102, 467
Castanea dentata, 28, 33, 35
Castilleja brunescens, 470; coccinea, 356; Haydeni, 99, 463, 464; lancifolia, 470; lauta, 470; miniata, 339; occidentalis, 100, 463; pallida, 99, 100, 102; rhexifolia, 470
Catabrosa aquatica, 474
Ceanothus americanus, 212; sanguineus 339
Celastrophyllum crenatum, 298; undulatum, 298
Celastrus arctica, 298
Centella repanda, 560
Cephalozia bicuspidata, 595; divaricata, 596; leucantha, 596, 615; media, 596; multiflora, 596
Cephaloziella byssacea, 596
Cerastrium alpinum, 94, 96; Behringianum, 96, 99, 467, 470; Earlei, 99, 468; pulchellum, 99, 468; vulgatum, 514, 516, 517, 518, 520
Ceratostoma, 161, 183, 184
Cercis canadensis, 214, 219, 243, 244, 251, 256, 533, 547, 548; Siliquastrum, 224
Cercis, On the relationship between the number of ovules formed and the number of seeds developing in, 243; Further observations, 533
Cercocarpus, 331, 497; alnifolius, 500; argenteus, 500; arizonicus, 502, 503; betulaefolius, 499, 500, 501; betulaefolius Blancheae, 500; betulaefolius typicus, 499; betuloides, 499, 500, 503; breviflorus, 501, 502; brevifolius, 501; brevifolius eximius, 501; Douglasii, 498, 499, 500, 503; flabellifolius, 500, 503; fothergilloides, 497, 498, 499; hypoleucus, 502, 503; intricatus, 502, 503; ledifolius, 502, 503; ledifolius intercedens subglabra, 502; ledifolius intricatus, 502; macrophyllus, 497; macrurus, 498, 503; minutiflorus, 500; mojadensis, 498; montanus, 498, 499, 500, 503; parvifolius, 498, 499, 500; parvifolius paucidentatus, 501; paucidentatus, 501; Pringlei, 498; rotundifolius, 499; Traskiae, 498, 500; Treleasii, 501
Cereus grandiflorus, 124
Chaenactis alpina, 464; Douglasii, 341; pedicularia, 464
Chaetomium, 183; fimeti, 182, 183
Chamaebatia, 331, 497; australis, 497
Chamaecyparis nootkatensis, 580; thyoides, 559, 563, 565
Chamaedaphne calyculata, 414
Chamaenerion angustifolium, 74, 75
Chamaepericlymenum canadense, 86
Cheirinia amoena, 100, 464; asperrima, 100; nivalis, 100, 464; Pallasii, 102; radiata, 100, 464; Wheeleri, 100
Chenopodium album, 514, 515, 516, 517, 518, 519; ambrosioides, 107, 127; hybridium, 512, 515
Chiloscyphus, 614; pallescens, 595; polyanthus, 614; polyanthus rivularis, 595; rivularis, 595
Chimaphila maculata, 28, 33, 411; umbellata, 308, 411, 415
Chiogenes hispidula, 426
Chionanthus virginica, 212, 214
Chloranthus, 232
Choiromyces, 182
Chondrophora nudata, 560, 563
Chondrophylla americana, 467; Fremontii, 469
Chondrosea Aizoon, 94, 466
Chrysanthemum Balsamitae tanacetoides,

514; Leucanthemum pinnatifidum, 515, 518, 519
Chrysoma laricifolia, 303
Chrysopsis alpicola, 464; aspera, 212; graminifolia, 561; mariana, 29, 35
Chrysosplenium alternifolium, 97; americanum, 28, 35; tetrandrum, 97, 468, 469
Cicuta bulbifera, 84; maculata, 84, 359
Cilia aggregata, 341
Cinchona Calisaya, 126; officinalis, 126; succirubra, 126
Cinnamomum Newberryi, 296, 297
Circaea alpina, 28, 32, 35; lutetiana, 82
Cirsium arvense, 518, 519; discolor, 355; Hillii, 356; lanceolatum, 515, 518, 519, 520
Cissites formosus magothiensis, 298; Newberryi, 298
Citrullus Colocynthis, 107, 126, 127
Cladium effusum, 560, 563
Cladonia sp., 212
Cladophlebis socialis, 296
Clavaria sp., 214 215
Claytonia linearis, 339; megarrhiza, 101, 465, 466; perfoliata, 339; sibirica, 339
Clematis hirsutissima, 340
Clementsia rhodantha, 470
Clerodendron calcicola, 12
Clethra alnifolia, 411, 414
Clethraceae, 411
Cliftonia monophylla, 560, 563, 567
Coccolobites cretaceus, 298
Coelopleurum actaeifolium, 85
Coffea, 149; arabica, 149
Colchicum autumnale, 107, 126.
Coleochaete, 192
Coleogyne, 331
Coleosanthus grandiflorus, 340
Collema, 164, 169, 179, 196; microphyllum, 165, 192; pulposum, 157, 159, 162, 163, 164, 166, 192, 197
Collinsia tenella, 339
Collinsonia canadensis, 109
Collomia grandiflora, 341
Colutea obovata, 297; primordialis, 297
Commandra umbellata, 354
Comocladia, 9; acuminata, 9; **cuneata,** 9; Dodonaea, 9; glabra, 9; pinnatifolia, 9; platyphylla, 9
Compsomyces verticillatus, 161
Condrophylla americana, 96; Fremontii, 99; humilis, 99; prostrata, 96
Conium maculatum, 83, 107; sp., 127
Conocephalum conicum, 585
Conopholis americana, 214, 215
Contributions to the Mesozoic flora of the Atlantic coastal plain, X—Maryland, 295
Convallaria, 225
Convolvulus Scammonia, 126; sepium, 355, 358, 360
Conyza squamata, 15
Copernicia, 20; **Cowellii,** 17; hospita, 20; macroglossa, 20; **rigida,** 17
Corallina, 178
Cordia apiculata, 298
Coreopsis major Oemleri, 212
Coriandrum sativum, 85, 107, 127
Cornaceae, 86
Cornus, 363; Amomum, 87, 362; canadensis, 86; cecilensis, 298; florida, 86, 214, 216, 219, 559, 563; Forchhammeri, 298; stolonifera, 28, 36, 338
Corylus americana, 362
Cowania, 495; alba, 495; Davidsonii, 495; ericaefolia, 495; Howardi, 495; mexicana, 495; Stansburiana, 495
Cracca virginiana, 212, 560, 562
Crataegus aestivalis, 566; brevispina, 341; Michauxii, 211, 214; sp. 214, 362
Crepis barbigera, 339; gracilis, 339
Crossosma Bigelovii, 303
Croton, 230; nummulariaefolius, 12
Crotonophyllum cretaceum, 297
Crucianella, 223, 225
Cruoriopsis, 196; cruciata, 175
Cudonia, 180
Cunilla originoides, 28, 33, 35
Cupressinoxylon Bibbinsi, 298
Cuscata californica, 341
Cynoglossum officinale, 514, 515, 518
Cynomorium, 222
Cynoxylon florida, 86
Cyperus comosus, 1; pseudovegetus, 561; sp. 358
Cypripedium, 229; acaule, 28, 32, 35, 36, 415; Reginae, 30
Cyrilla racemiflora, 560, 561, 567
Cytisus scoparius, 107, 124
Czekanowskia capillaris, 296

Dalbergia severnensis, 297
Dammara borealis, 299; cliffwoodensis, 299
Danthonia intermedia, 469
Dasiophora fruticosa, 30, 463
Dasylirion Wheeleri, 302, 303
Dasystephana affinis, 463; frigida, 96; Froelichii, 96; glauca, 95, 468; monticola, 464; Parryi, 464; Romanzovii, 96
Dasystoma laevigata, 28, 33, 35; quercifolia, 214
Datura, 121; ferox, 127; Stramonium, 107, 121, 444; Tatula, 121
Daucus Carota, 86, 360
Decodon verticillatus, 73
Delphinium alpestre, 101, 468; Consolida, 107, 121; Menziesii, 340; Stephisagria, 107, 127
Dendropanax, 9; cuneifolium, 10; **filipes,** 9
Dentaria heterophylla, 28, 35
Denticulatae, 591, 592
Dermonema, 194; dichotomum, 172

Deschampsia alpicola, 98, 469, 470; caespitosa, 93, 96, 98, 469; curtifolia, 96, 463
Desmodium illinoense, 355
Dianthus barbatus, 516
Dichromena latifolia, 560, 566
Digitalis, 115; alba, 116, 122, 123; ambigua, 116, 122, 123; Buxbaumi, 122, 123; canariensis, 116, 122, 123, 124; gloxiniaeflora, 116, 122, 123; gloxiniaeflora alba, 117; gloxiniaeflora rosea, 116; grandiflora, 122, 123; lanata, 116, 122, 123; lutea, 122, 123; macrantha, 122, 123; maculata Iveryana, 116; monstrosa, 122, 123; purpurea, 116, 122; purpurea alba, 122, 123; purpurea maculata superba, 122; purpurea rosea, 122, 123; sibirica, 116, 122, 123
Dilophozia, 589
Dioscorea sp., 214
Diospyros primaeva, 297, 298; rotundifolia, 298; vera, 297; virginiana, 28, 34, 559, 560
Diplophylleia albicans, 602; plicata 602; taxifolia, 602
Diplophyllum albicans, 590, 602, 615; argenteum, 613, 615; imbricatum, 613, 615; ovatum, 602, 615; plicatum, 602, 614; taxifolium, 602
Disporum majus, 339
Dodecatheon Meadia, 29, 32, 35, 356, 357; puberulum, 339
DODGE, B. O. The morphological relationships of the Florideae and the Ascomycetes, 157
Dodonaea angustifolia, 303
Doryanthites cretacea, 298
Douglas Lake, Michigan, The introduced vegetation in the vicinity of, 511
Douglasia montana, 101, 463, 464; nivalis, 101, 465
Draba, 462; alpina, 102; andina, 96, 99, 462, 463; aurea, 463; aureiformis, 464; chrysantha, 464; crassa, 464; crassifolia, 464; densiflora, 462, 463; fladnizensis, 93, 467; glacialis, 94, 96; lonchocarpa, 98, 468; nivalis, 98, 102, 468; oligosperma, 96, 462, 463; Parryi, 468; pectinata, 99; saximontana, 464; streptocarpa, 464, 470
Drosera filiformis Tracyi, 560, 563, 567
Dryas, 483; Drummondii, 462, 464, 483, 484; integrifolia, 483; integrifolia canescens, 483; integrifolia intermedia, 483; octopetala, 462, 463, 483, 484, 490; octopetala intermedia, 483; octopetala argentea, 483, 484; octopetala hirsuta, 483; octopetala minor, 483; tomentosa, 483
Drymocallis agrimonioides, 485; Convallaria, 339; glandulosa, 101; pseudorupestris, 99, 471; pumila, 101; rupestris, 99
Drymoglossum, 397
Dudresnaya, 167, 176, 177, 194, 195, 196; coccinea, 173, 195; purpurifera, 195

Eatonia nitida, 30, 35
Eichhornia, 225
Elaeodendron marylandicum, 298
Elaphomyces, 182
Elephantella, 472; groenlandica, 471, 472
Elettaria Cardamomum, 107, 127
Elymus canadense, 356
Endogone, 157, 192
Epigaea repens, 28, 33, 35, 414
Epilobium adenocaulon, 76; angustifolium, 517, 519; coloratum, 76; hirsutum, 75; lineare, 75, 76; palustre, 75; palustre labradoricum, 75; palustre monticola, 75; strictum, 76
Epipactis, 143, 223, 224, 225, 231; pubescens, 230
Equisetum arvense, 356
Erechtites hieracifolia, 517
Erica cinerea, 418; Tetralix, 416, 417
Ericaceae, 413
Erifonum flavum, 100; xanthum, 100
Erigeron alpinum, 94; annuus, 360; canadensis, 360, 517; compositus, 97, 462, 463; expansus, 15; glabellus, 470; grandiflorus, 102; lanatus, 102; melanocephalus, 96; multifidus, 97, 99, 462, 463; pedatus, 99; pinnatisectus, 462, 464; ramosus, 354, 356; simplex, 96, 462, 463; speciosus, 339; trifidus, 97, 462, 463; unalaschensis, 102; uniflorus, 96
Eriocaulon decangulare, 560, 566, 567
Eriogonum compositum, 341; flavum, 100; heracleoides, 341; niveum, 341, 342; tomentosum, 563; xanthum, 100, 464
Eriophorum gracile, 471; polystachyum, 471
Eriophyllum lanatum, 341, 342
Eritrichium argenteum, 99, 462, 464; elongatum, 99, 463, 464; nanum, 99; villosum, 99
Ernodea, 12; littoralis, 12
Eryngium, 359; yuccifolium, 356, 357, 359
Erysimum asperum, 340
Erythrina Crista-galli, 224; flabelliformis, 303
Erythrocoma, 490, 491, 492; affinis, 492; aliena, 492; arizonica, 492; australis, 492; canescens, 493; ciliata, 490; cinerascens, 492, 493; companulata, 491; grisea, 492; tridentata, 492
Erythronium grandiflorum, 339
Erythrophyllum, 173, 178, 194, 197; delesserioides, 173
Erythroxylum Coca, 107, 126
Eucalyptus attenuata, 298; Geinitzi, 298; latifolia, 298; Wardiana, 298

Eujungermannia, 588
Eupatorium aromaticum, 28, 32, 35; capillifolium, 560, 563; coelestinum, 28, 33, 35; compositifolium, 560, 563; perfoliatum, 358, 359; purpureum, 560; rotundifolium, 28, 32, 35, 560
Euphorbia, 223, 230; corollata, 355, 357; cyparissias, 514; hirsuta, 512; maculata, 512; palustris, 224, 230; procera, 224, 230; virgata, 229
Euthamia grandiflora, 340
Euvittaria, 394, 395
Euxolus emarginatus, 509
EVANS, A. W. Report on the Hepaticae of Alaska, 577
Exoascus, 170

Fagus grandifolia, 213, 216, 219, 512, 559
Fallugia, 496; acuminata, 496; mexicana, 496; micrantha, 496; paradoxa, 496
FARWELL, O. A. The correct name for the hemlock spruce, 621
Fatsia japonica, 222
Fegatella conica, 585
Ferns and flowering plants of Nantucket, The—XII, 71;—XIII, 411
Festuca brachyphylla, 99, 463; ingrata, 469; minutiflora, 99, 464; ovina, 93, 96; rubra, 93, 469; saximontana, 96, 463, 469; supina, 93, 463, 464; Thurberi, 469, 470
Ficus Cecilensis, 298; crassipes, 298; daphnogenoides, 298; Krausiana, 298; ovatifolia, 297
Fimbriaria tenella, 613
FISHER, G. C. Seed development in the genus Peperomia, 137, 221
Foeniculum vulgare, 107, 126, 127
Fontainea grandifolia, 297
Fragaria platypetala, 339; virginiana, 356
Fraxinus americana, 213, 219; nigra, 29, 36; quadrangulata, 28, 36
Frullania chilcootiensis, 613, 615; franciscana, 612, 615; nisquallensis, 590, 607, 612, 614, 615; Tamarisci, 614
Fucus, 229

Gaillardia aristata, 342
Galactinia, 158; succosa, 170
Galega officinalis, 107
Galium Aparine, 339; boreale, 339; sp. 357; uniflorum, 214
Garcinia, 222
Gaultheria procumbens, 28, 33, 34, 36, 414
Gaylussacia baccata, 419; dumosa, 212, 419, 421; dumosa Bigeloviana, 419; frondosa, 419
Geinitzia formosa, 298
Gelsemium sempervirens, 212
Geocalyx graveolens, 595
Geoglossum, 171, 180
Gentiana Andrewsii, 359; crinita, 29, 36; lutea, 126; oregana, 339
Geranium viscossisimum, 339
Geum, 484, 490, 491, 494; agrimonioides, 484, 485; album, 485, 486; album flavum, 486; aleppicum, 488; auranthacum, 489; calthifolium, 494; camporum, 484; canadense, 484, 485, 486; cercocarpoides, 496; ciliatum, 491; decurrens, 486; flavum, 486; florum, 486; geniculatum, 488; Grahami, 492; hirsutum, 486; intermedium, 488; laciniatum, 484; macrophyllum, 487, 488; macrophyllum × oregonense, 489; macrophyllum × rivale, 489; macrophyllum × strictum, 489; mexicanum, 488; Meyerianum, 484, 485, 486; nutans, 488; oregonense, 487; oregonense × strictum, 489; perincisus, 487; pulchrum, 489; Rafinesquianum, 488; rivale, 484, 488, 489, 490; rivale × strictum, 489; rotundifolium, 494; scopulorum, 487; strictum, 363, 486, 487, 488, 489; triflorum, 491; urbanum, 487, 488, 489; vernum, 484; virginianum, 484
Gigartina, 173, 178
Ginkgo biloba, 90
GLEASON, H. A., and MCFARLAND, F. T. The introduced vegetation in the vicinity of Douglas Lake, Michigan, 511
Gleichenia delawarensis, 297; Saundersii, 297; Zippei, 297
Gloiosiphonia, 176, 177, 194
Glyceria nervata, 358, 359, 360
Glycyrrhiza glabra, 126
Gnomonia, 158, 171; erythrostoma, 159
Gonianthes Lindeniana, 23; Sagraeana, 23
Gormania debilis, 101, 464
Graphis elegans, 164
Griffithsia Bornetiana, 167
GRIGGS, R. F. Observations on the behavior of some species at the edges of their ranges, 25; Observations on the edge of the forest in the Kodiak region of Alaska, 381
Grimaldia fragrans, 612
Grindelia robusta, 127
Grossularia hirtella, 415
Gunnera, 137, 138, 223; macrophylla, 137
Gymnadenia, 149, 150; conopsea, 149
Gymnoascus, 171, 193; candida, 159
Gymnocladus dioica, 29, 35
Gymnomitrium concinnatum, 612; corallioides, 612; crenulatum, 612, 615; obtusum, 586, 615
Gyrostachys gracilis, 33
Gyrothyra Underwoodiana, 587, 615

Habenaria nivea, 561
Halesia diptera, 219
Haloragidaceae, 82
Hamamelis virginiana, 214
Haplopappus marginatus, 14
Haplopteris, 393, 394, 395

HARPER, R. M. The "Pocosin" of Pike County, Alabama, and its bearing on certain problems of succession, 209; A superficial study of the pine-barren vegetation of Mississippi, 551
Harrimanella hypnoides, 467
HARRIS, J. A. On the relationship between the number of ovules formed and the number of seeds developing in Cercis, 243; Further observations, 533
Harveyella, 173, 174, 194, 197; mirabilis, 167, 172
Havardia, 8
Hecistopteris, 400; minima, 398; Weckleana, 399
Hedera cecilensis, 298; cretacea, 298; Helix, 367, 368, 370, 371
Hedyosmum, 232
Helenium tenuifolium, 560
Helianthus, 363; angustifolius, 561; annuus, 512; decapetalus, 362; divaricatus, 362; grosse-serratus, 358, 360; Maximiliani, 355; occidentalis, 355; scaberrimus, 355
Heliopsis scabra, 355
Helminthora, 167
Helotium citrinum, 190
Helvella, 179
Hemlock spruce, The correct name for the, 621
Hepaticae of Alaska, Report on the, 577
Heracleum lanatum, 85, 86, 339
Herberta adunca, 597, 601, 615
Heteromallae, 592
Heuchera cylindrica, 339; Hallii, 465; hispida, 356; parvifolia, 463
Hibiscus aculeatus, 561
Hicoria alba, 216, 219, 559; glabra, 211, 213
Hieracium albiflorum, 339; paniculatum, 28, 35; Scouleri, 339; venosum, 28, 33, 35
Hoffmannia aphylla, 56
Hordeum jubatum, 512
Houttuynia, 232
HOWE, R. H., JR. The nomenclature of the genus Usnea, 373
Hulsea carnosa, 465, 466
Humaria granulata, 159; rutilans, 171
Humulus Lupulus, 512
Hydrangea arborescens, 28, 35
Hydrastis canadensis, 109
Hydnocystis, 182
Hydrocotyle umbellata, 83
Hyoscyamus muticus, 126; niger, 107, 125, 444; orientalis, 125
Hypericum Drummondii, 29, 32, 35; fasciculatum, 560; sp. 359
Hypocopra, 161, 183
Hypopitys americana, 412; **insignata,** 413; lanuginosa, 412
Hypoxis hirsuta, 358

Ichmadophila aeruginosa, 159
Ilex coriacea, 216; fastigiata, 414; glabra, 559, 567; myrtifolia, 560, 563, 567; opaca, 214, 216, 219, 559; severnensis, 298
Illicium, 148; deletoides, 298
Impatiens biflora, 339
Inga Hystrix, 4; obovalis, 5
Introduced vegetation in the vicinity of Douglas Lake, Michigan, The, 511
Ionactis linariifolius, 212
Iris, 149; cristata, 29, 35, 36; sibirica, 149; verna, 212; versicolor, 359
Isnardia palustris, 74
Isoetes Bolanderi, 474; paupercula, 474
Isopyrum biternatum, 28, 35
Isotria verticillata, 29, 34
Iva xanthifolia, 512, 514

Jacobinia **jamaicensis,** 16
Jeffersonia diphylla, 109
Juglans, 231
Juncoides arcticum, 102, 471; arcuatum, 102, 471; hyperboreum, 102, 471; parviflorum, 470; saltuensis, 28, 32, 35; spicatum, 93, 94, 467, 470
Juncus, 466; biglumis, 102, 471; castaneus, 93, 467, 470; Drummondii, 467, 470; Parryi, 467, 470; Roemerianus, 560, 563; sp. 357; triglumis, 93, 467, 470
Jungermannia, 588; albicans, 602; albicans taxifolia, 602; atrovirens, 587; caespiticia, 613, 615; cochleariformis, 610; conchaefolia, 611; cordifolia, 587; gigantea, 611; lanceolata, 613; minuta, 588; nemorosa, 606; orcadensis, 588; purpurea, 610; saxicola, 613; setiformis, 613; sphaerocarpa, 613; trichophylla, 601; ventricosa, 613
Jungermanniaceae, 586, 612
Juniperus australis, 11; hypnoides, 298; lucayana, 11; scopulorum, 343

Kalmia, 467; angustifolia, 414; glauca, 97; latifolia, 28, 33, 35, 214, 216, 413; microphylla, 97, 472, 473
Kantia Trichomanis, 596
Kerria, 331; japonica, 367, 370
Kieseria, 20; **cubensis,** 19
Kneiffia pumila, 82
Koeleria cristata, 354, 356
Koellia incana, 28, 35
Krigia amplexicaulis, 356
Kuhnistera pinnata, 212, 563
KUNKEL, L. O. Physical and chemical factors influencing the toxicity of inorganic salts to Monilia sitophila (Mont.) Sacc., 265
Kuntzia, 496, 497; tridentata, 495

Laboulbenia chaetophora, 159
Lachemilla, 319, 320, 321; Bourgeaui,

321; domingensis, 321; ocreata, 321; orbiculata, 320, 321; orizabensis, 321; Pringlei, 321; procumbens, 321; Schiedeana, 321; venusta, 320
Lachnea, 169, 180, 186; abundans, 163, 165, 189, 193; cretea, 159, 160, 165, 167, 172, 189, 193; leucotricha, 189; melaloma, 166, 193; scutellata, 159, 165, 171, 181, 185, 186, 189, 193, 194; stercorea, 159, 164, 181, 185, 186, 189, 192, 193
Lacistema, 232
Lactuca canadensis, 360
Lappula ciliata, 341; deflexa, 512, 514, 515, 518
Larix, 153; sibirica, 222
Lasiobolus, 195; equinus, 159, 160, 193, 194; pulcherrimus, 193, 194
Lasiocroton Fawcettii, 16; **Harrisii**, 16
Lathyrus palustris, 356
Laurus Cinnamomum, 126; Hollickii, 297; plutonia, 297; proteaefolia, 297
Laurophyllum angustifolium, 297; elegans, 297
Lecanora subfusca, 166
Lechea racemulosa, 28, 36
Ledum glandulosum, 467, 472, 473
Leguminosites canavalioides, 297; coronilloides, 297; omphalobioides, 297
Lejolisia, 160
Lemna, 231
Leotia, 171; chlorocephala, 159
Lepachys pinnata, 355, 357, 360
Lepidium apetalum, 512; virginicum, 512, 514, 515, 516, 517, 518, 520
Lepidozia, 599, 600; filamentosa, 598, 600, 615; Pearsoni, 601; reptans, 598, 600; sandvicensis, 598, 599, 600, 601, 614, 615; setacea, 601, 614
Leptandra virginica, 109
Leptasea austromontana, 96, 466; bronchialis, 96, 99; chrysantha, 98, 468; flagellaris, 467; Hirculus, 98, 468, 469, Van Bruntiae, 99, 469
Leptilon canadense, 561
Leptinae, 377
Lespedeza capitata, 355, 360
Levisticum officinale, 126
Liagora, 167
Liatris scariosa, 354, 361; spicata, 358, 359, 360
Lichen articulatus, 376, 377; barbatus, 376, 377; floridus, 374; hirtus, 374, 375; plicatus, 375
Ligusticum Porteri, 472; scoticum, 85
Lilium, 229, 231; canadense, 356
Linum floridanum, 561; intercursum, 415
Liquidambar Styraciflua, 219, 559, 563
Liriodendron Tulipifera, 41, 46, 216, 219, 559
Liriodendropsis constricta, 297
Lithospermum canescens, 355; ruderale, 341
Lloydia serotina, 90, 93, 95, 470
Lobelia inflata, 107, 127; leptostachys, 29, 36; puberula, 29, 35; spicata, 356; syphilitica, 359
Lomatium Grayi, 341; triternatum, 340
Lonicera japonica, 367
Lophiola aurea, 560, 563, 566
Lophocolea cuspidata, 595, 615
Lophozia, 588, 589; attenuata, 613; Floerkei, 587; guttulata, 613; heterocolpa, 613; incisa, 587; inflata, 587; lycopodioides, 613; minuta, 588; obtusa, 613, 615; porphyroleuca, 587; quadriloba, 613; quinquedentata, 588; ventricosa, 613
Ludwigia alternifolia, 74; pilosa, 561
Lupinus albus, 267; microcarpus, 343; Piperi, 341
Lychnis alba, 514, 518
Lycium carolinianum, 12; halimifolium, 367, 368, 369
Lyopodiales, 51
Lycopodiaceae, 51
Lycopodium, 51, 55; alopecuroides, 561; arbuscula, 58; capillaceum, 54; cernuum, 51, 53; clavatum, 29, 32, 35, 36, 38, 51; compactum, 53; complanatum, 29, 33, 35, 36, 37; crassifolium, 54; curvatum, 53; digitatum, 57; erubescens, 51, 52, 53; Haleakalae, 51, 53; javanicum, 52; lucidulum, 28, 33, 35, 36, lucidulum porophilum, 28, 34, 36; macrostachys, 55; Menziesii, 58; nudum, 56; nutans, 51, 54; obscurum, 29, 35, 36, 38; pachystachyon, 54; pachystachyon phyllanthon, 54; pennigerum, 58; Phlegmaria, 51, 53; phyllanthum, 51, 54, 55; polytrichoides, 52, 56; Sabinae-facie, 57; sargassifolium, 52; selaginoides, 57; serratum, 51, 52, 58; serratum dentatum, 52; serratum subintegrum, 52; squarrosum, 54; suberectum, 53; sulcinervium, 52; varium, 52; venustulum, 52, 55; venustulum herpeticum, 55; volubile, 52, 55
Lysias orbiculata, 30
Lysimachia quadrifolia, 28, 34
Lythraceae, 73
Lythrum Salicaria, 73; Salicaria tomentosum, 73

Machaeranthera canescens, 342; Pattersoni, 465
Macronema Lyallii, 465; pygmeum, 464
Madotheca navicularis, 611; platyphylla, 613
Magnolia acuminata, 219; Boulayana, 298; capellinii, 298; glauca, 216, 559;

grandiflora, 213, 216, 219, 559, 563; Hollicki, 298; Lacoeana, 298; longipes, 298; macrophylla, 219, 559, 563, 566; obtusata, 298; pyramidata, 219; tenuifolia, 298
Mairania alpina, 102
Malva rotundifolia, 514, 516
Marchantia, 333; polymorpha, 585, 615
Marchantiaceae, 585
Marsupella emarginata, 586; sphacelata, 612
Maryland, Contributions to the Mesozoic flora of the Atlantic coastal plain, —X., 295
Mastigobryum Didrichsenii, 597; Pearsoni, 596; deflexum, 598
Maytenus phyllanthoides, 12
McFARLAND, F. T., GLEASON, H. A., and, Introduced vegetation in the vicinity of Douglas Lake, Michigan, The, 511
Mechanical tissue development in certain North American vines, 365
Medicinal plants, The propagation of, 105
Meibomia angustifolia, 19; **Cowellii**, 19
Melampyrum lineare, 29, 32, 35
Melanospora, 161, 183, 184
Melastomaceae, 74
Melilotus alba, 360
Mentha canadensis borealis, 339
Mentzelia integrifolia, 342; laevicaulis, 342
Mertensia alpina, 466, 467, 471; Bakeri, 100, 468; brevistyla, 468, 471; Drummondii, 102; lanceolata, 100; lateriflora, 100, 468; nivalis, 100; oblongifolia, 339; Parryi, 100, 468; viridula, 100, 468
Mesinae, 374
Mesozoic flora of the Atlantic coastal plain, Contributions to the, X.—Maryland, 295
Mesophylla orcadensis, 588
Metzgeriaceae, 585
Metzgeria conjugata 586, 590, 593, 607; hamata, 612, 614, 615; pubescens, 586
Micranthes arnoglosa, 98, 468; brachypus, 468; crenatifolia, 98, 469; foliolosa, 102; hieracifolia, 94, 98, 469; nivalis, 94, 98, 102; rhomboidea, 98, 467; Rydbergii, 98, 468; stellaris, 468; Vreelandii, 468, 469
Microglossum, 180
Microlepidozia, 614
MILLER, F. A. The propagation of medicinal plants, 105
Mimosa arborea, 7; biuncifera, 303; circinalis, 3; filicifolia, 7; Saman, 8; Unguis-cati, 3
Mitchella repens, 214
Mitrula, 180
Monarda, 109; mollis, 356, 363
Monilia sitophila (Mont.) Sacc., Physical and chemical factors influencing the toxicity of inorganic salts to, 265
Monilia sitophila, 266, 268, 269, 272, 275, 277, 281, 283, 284, 286, 289
Monotropaceae, 412
Monotropa, 149, 150; Hypopitys, 149; uniflora, 214, 215, 412
Morchella, 169
Moriconia americana, 298
Morphological relationships of the Florideae and the Ascomycetes, The, 157
Morus, 148, 149, 150; alba, 146, 147, 148
Mosaic disease of tobacco, A review of investigations of the, together with a bibliography of the more important contributions, 435
Mucor, 179; Mucedo, 286, 290
Muscaria adscendens, 93, 467; caespitosa, 98, 102, 468; delicatula, 98, 468; micropetala, 98, 468; monticola, 98, 469
Muscus clavatus, 57
Mylia anomala, 595; Taylori, 595
Myosotis alpestris, 93, 466, 467
Myrica cerifera, 560; inodora, 566; longa, 298; pumila, 560
Myriophyllum humile, 82; tenellum, 82
Myrsine borealis, 298; Gaudini, 298

Nantucket, The ferns and flowering plants of,—XII, 71; XIII, 411
Napaea dioica, 29, 34, 38
Nardia compressa, 613, 615; Geoscyphus, 613, 615; haematosticta, 613; obovata, 613, 615; scalaris, 587
Nelumbites primaeva, 297
Nemalion, 167, 171, 172, 174, 194, 195
Nepeta Cataria, 514, 515, 516, 518, 519
New or notable species of Amaranthus, 505
Nicandra physaloides, 125
Nomenclature of the genus Usnea, The, 373
North American ferns, Notes on two, 307
Notes on Rosaceae—VII, 319; VIII, 483
Notes on two North American ferns, 307
Nyssa biflora, 559, 560; Ogeche, 563; sylvatica, 87, 211, 219, 559, 563; uniflora, 559, 563

Observations on the behavior of some species at the edges of their ranges, 25
Observations on the edge of the forest in the Kodiak region of Alaska, 381
Oedocephalum albidum, 290
Oedogonium, 167
Oenothera, 229, 366, 371; albida, 81; biennis, 76, 77, 78, 80, 355; biennis parviflora, 79; canovirens, 78; cruciata, 80; Lamarckiana, 80, 81, 366; muricata,

76, 77, 80; Oakesiana, 78, 80; rubrinervis, 366; **stenopetala**, 79, 80; strigosa, 77; subulifera, 78
Oetosis, 395, 396, 397
Onagraceae, 74
Onoclea inquirenda, 297; sensibilis, 618; Struthiopteris, 617
Onoclea sensibilis L., Branched cells in the prothallium of, 617
Onosmodium, 497
On the relationship between the number of ovules formed and the number of seeds developing in Cercis, 243; Further observations, 533
Opulaster pauciflorus, 339
Opuntia, 71, 72; Opuntia, 72; polyacantha borealis, 343; vulgaris, 72, 212, 214
Orchis, 150; latifolia, 149; Morio, 149
Oreobroma pygmaea, 463
Oreocarya sericea, 342
Oreochrysum Parryi, 468, 471
Oreoxis alpina, 101, 464; Bakeri, 101, 464; humilis, 101, 464
Origin of the alpine flora. Phytogeographical notes on the Rocky Mountain region, II, 89
Ornithogalum, 229
Osmanthus americana, 212, 214, 216
Osmunda cinnamomea, 561; delawarensis, 297
Ostrya virginiana, 214, 219
Oxalis stricta, 358
Oxycoccus, 472; macrocarpus, 426; Oxycoccos, 426
Oxydendrum arboreum, 28, 33, 35, 37, 214, 219, 559, 562, 563
Oxypolis filiformis, 560, 567; rigidior, 356, 359, 363
Oxyria digyna, 93, 465, 466, 467

Pallavicinia Flotowiana, 586, 615; hiberica, 586
Panicularia acutiflora, 30; elongata, 30, 36; palida, 30
Panicum capillare, 512; dichotomum, 561; nitidum, 357; polyanthes, 28, 35; Scribnerianum, 356; sp., 214, 357; stipitatum, 30, 35; virgatum, 354
Papaver somniferum, 107
Parnassia caroliniana, 29, 36
Paronychia pulvinata, 101, 464
Parrya macrocarpa, 98; platycarpa, 98, 464
Parthenium integrifolium, 356, 357, 359
Passiflora incarnata, 107; lutea, 29, 34, 38
Pastinaca sativa, 85
Pedicularis canadensis, 356; flammea, 102; lanata, 102; lanceolata, 29, 35; Oederi, 102; Parryi, 470; racemosa, 470
Pellaea ternifolia, 308
Pellia endiviaefolia, 612; epiphylla, 612; Fabroniana, 612; Neesiana, 586
Penaea, 223
Penicillium, 170, 173, 178; crustaceum, 289; glaucum, 290
Pentstemon confertus, 339; Hallii, 464, 465; Harbourii, 465; stenosepalum, 468
Peperomia, Seed development in the genus, 137, 221
Peperomia, 137, 138, 140, 141, 146, 148, 150, 151, 152, 154, 156, 221, 222, 223, 224, 225, 226, 229, 230, 231, 232, 233, 234; arifolia, 140, 143, 223, 224; blanda, 138, 139, 154, 225; Fraseri, 153; Fraseri resediflora, 138, 139, 153, 225; galioides, 138, 139, 155, 225; hispidula, 137, 142, 223; humilis, 139; ionophylla, 140; Langsdorffii, 138, 139, 155, 225; metallica, 138, 139, 152, 153; Ottoniana, 223; pellucida, 137, 138, 142, 144, 223; pulchella, 145; reflexa, 138, 139, 145, 146, 150, 151, 152, 154, 155, 225; resedaeflora, 139, 153; scandens, 138, 150, 151, 225; Sintenisii, 143, 223, 224; urocarpa, 140; verticillata, 138, 139, 145, 147, 148, 149, 151, 152, 154, 225
Peronospora, 160
Persea Borbonia, 214
Pertusaria communis, 159
Petalostemum purpureum, 355
Petrocelis, 176
Petroselinum hortense, 84
Petunia nyctaginifolia, 444
Phacelia alpina, 100, 464, 471; ciliosa, 470; dubia, 29, 32, 35; heterophylla, 100, 342; Lyallii, 99, 465; sericea, 99, 464, 470
Phallus, 179
Phaseolus, 256, 548; multiflorus, 224; vulgaris, 243, 548
Philadelphus Lewisii, 341
Phippsia algida, 97, 474
Phleum alpinum, 93, 94, 469, 471; pratense, 360, 514, 515, 516, 518, 519, 520
Phlox, 462; caespitosa, 100, 462, 463; condensata, 100, 462, 464; glaberrima, 356; pilosa, 356, 358; stolonifera, 29, 32, 34, 36, 38
Phoradendron flavescens, 560
Phragmites communis, 359
Phycomyces nitens, 286
Phyllactinia corylea, 159
Phyllanthus, 230
Phyllodoce, 467; coerulea, 98; empetriformis, 98, 472, 473; glanduliflora, 98, 472, 473
Physalis virginiana, 354
Physaria Geyeri, 342
Physical and chemical factors influencing the toxicity of inorganic salts to Monilia sitophila (Mont.) Sacc., 265
Physiotium cochleariforme, 610
Physostegia virginiana, 358

Physma compactum, 192
Phytogeographical notes on the Rocky Mountain region, II. Origin of the alpine flora, 89; III. Formations in the alpine zone, 459
Picea alba, 628; balsamea, 628; canadensis, 621, 628, 629; mariana, 628; nigra, 628; sitchensis, 381, 580
Pimpinella Anisum, 107, 127
Pine-barren vegetation of Mississippi, A, superficial study of, 551
Pinus, 153; alba, 628; americana, 628; balsamea, 623, 624, 625, 626, 627, 628; canadensis, 621, 622, 623, 625, 626, 627, 628, 629; contorta, 343; echinata, 211, 559, 563; Elliottii, 559, 563, 564; glabra, 219, 559, 563, 565; mariana, 628; nigra, 628; palustris, 211, 551, 559, 561; ponderosa, 342; rigida, 29, 32, 33, 35; serotina, 209, 563; Taeda, 219, 559, 563, 565; virginiana, 29, 33, 35
Piper, 137, 138, 140, 141, 153, 155, 156, 229, 231, 232; Betle monoicum, 156; medium, 156; nigrum, 126; tuberculatum, 138, 155, 156, 233; umbellata, 156
Pistia Nordenskioldi, 298
Pitcheria galactioides, 212
Pithecolobium, 2; arboreum, 3, 7; asplenifolium, 3, 6; bahamense, 2, 4; Berterianum, 3, 8; brevifolium, 8; calliandrifolium, 4; circinale, 2, 3; **discolor,** 2, 4; filicifolium, 7; guadalupense, 2, 4; **guantanamense,** 3, 7; Hystrix, 2, 4; latifolium, 8; mucronatum, 3; **nipense,** 3, 6; obovale, 2, 5; **pinetorum,** 2, 6; prehensile, 3, 8; Saman, 3, 8; **savannarum,** 2, 4; tortum, 3, 8; **trinitense,** 3, 7; **truncatum,** 2, 5; Unguis-cati, 2, 3; vincentis, 8
Plagiochila, 589, 591, 594; **alaskana,** 590, 591, 592, 594, 595, 615; asplenioides, 590; fragilis, 592, 593; **Fryei,** 593, 594, 595, 615; oxyphylla, 592, 593; pachyloma, 592, 593; renitens, 591, 592; trapezoidea, 591, 592
Plantago lanceolata, 514, 515; major, 512, 514, 515, 516, 518, 519
Platanus, 297; Heerii, 297
Pleuroschisma Pearsoni, 596
Pleurozia, 610; cochleariformis, 610; conchaefolia, 611; gigantea, 611; purpurea, 610, 611, 614, 615, 616
Poa, 469; alpicola, 470; alpina, 93, 94, 467; annua, 515; arctica, 96, 99, 467; Buckleyana, 469; cenisia, 93, 96, 467; compressa, 356, 357, 363, 514, 515, 516, 518, 520; crocata, 100, 101, 463; epilis, 464, 469; Grayana, 470; leptocoma, 469, 471; Lettermannii, 101, 463, 464; longiligula, 469; Pattersonii, 100, 463, 464; pratensis, 360, 512, 514, 515, 516, 518, 519, 520, 521; pudica, 99, 470; reflexa, 471; rupicola, 101, 463; Sheldoni, 470
"Pocosin" of Pike County, Alabama, and its bearing on certain problems of succession, The, 209
Podostemon, 229
Podozamites Knowltoni, 297; lanceolatus, 296; marginatus, 296
Polemonium, 466; Brandegei, 101, 468; confertum, 101, 465, 468, 470; delicatum, 99, 466; Grayanum, 101; humile, 96; mellitum, 101, 470; micranthum, 343; parviflorum, 99, 469; pulcherrimum, 96, 99, 466, 467; speciosum, 101, 465, 470; viscosum, 101, 466, 468, 471
Polygala cruciata, 561; cymosa, 560, 566; ramosa, 560, 567
Polygonatum biflorum, 214
Polygonum amphibium Hartwrightii, 360; arifolium, 32, 35; aviculare, 512, 514, 515, 518, 520; convolvulus, 514, 515, 517, 518, 519; erectum, 512; Persicaria, 514, 515, 518; sp. 356
Polyides, 167, 173, 176
Polypodium, 391; polypodioides, 212, 214
Polysiphonia, 167, 174
Polystigma, 158, 171, 191
Populus balsamifera, 381, 580; deltoides, 359; grandidentata, 511; stygia, 298; tremuloides, 29, 36; trichocarpa, 338, 340
Porella, 608; navicularis, 611, 615; platyphylla, 613; rivularis, 612; Roellii, 612, 615
Poronia punctata, 159
Porteranthus stipulatus, 29, 35
Portlandia, 21; coccinea, 21, 23; coriacea, 23; daphnoides, 21, 23, 24; **domingensis,** 21, 24; **elliptica,** 21, 22; grandiflora, 21, 22; gypsophila, 23; Harrisii, 21, 22; involucrata, 21, 22; **Lindeniana,** 21, 23; longiflora, 23; nitens, 21, 22; pendula, 21, 23; **sessilifolia,** 21; uliginosa, 21, 22
Portulaca oleracea, 517
Potamogeton alpinus, 474
Potentilla, 92, 495; arguta, 485; decurrens, 100; diversifolia, 100, 470; emarginata, 102; glaucophylla, 470; Macounii, 464; minutiflora, 101; minutifolia, 464; modesta, 98; nipharga, 96, 464; nivea, 90, 92, 93, 96, 98, 464; paucijuga, 464; perdissecta, 100; quinquefolia, 96, 98, 462, 463; Robbinsiana, 92; rubripes, 470; saximontana, 101; tenerrima, 101, 464; uniflora, 93, 102; villosa, 93, 102
Poteridium, 322; annuum, 322, 323; occidentalis, 323
Poterium, 323; annuum, 323; Sanguisorba, 323; spinosum, 323

Preissia hemisphaerica, 585; quadrata, 585
Primula angustifolia, 100, 467, 470; Broadheadae, 100; farinosa, 90; Parryi, 465, 470, 472
Prionitis, 194, 197; Lyallii, 174
Propagation of medicinal plants, The, 105
Proserpinaca palustris, 82
Prosopis velutina, 303
Protophyllocladus lobatus, 297; subintegrifolius, 297
Protophyllum, 296, 297; multinerve, 297; Sternbergii, 297
Prunus americana, 362; caroliniana, 214; demissa, 340; serotina, 213; umbellata, 214
Psedera quinquefolia, 363, 366, 369
Pseudocymopterus montanus, 100; purpureus, 471; Tidestromii, 100, 464
Pseudopteryxia anisata, 465; Hendersonii, 466
Pseudotsuga taxifolia in arid parts of Washington, Slope exposure as a factor in the distribution of, 337
Pseudotsuga taxifolia, 337, 338, 339, 340, 343, 344
Psilotaceae, 56
Psilotum, 56; complanatum, 56, 57; nudum, 56; oahuensis, 56; triquetrum, 56
Psoralea Onobrychis, 29, 36
Psychotria, 11; ligustrifolia, 11; pulverulenta, 11; Sulzneri, 11; undata, 11
Pteridium aquilinum, 212
Pteridophyta of the Hawaiian Islands, A taxonomic study of the,—IV, 51
Pteris, 395, 396, 397; lanceolata, 397; lineata, 393, 394, 395, 397; piloselloides, 397; scolopendrina, 394
Pteropsis, 397
Ptilidium californicum, 601, 615; ciliare, 602; pulcherrimum, 602
Ptilimnium capillaceum, 84
Ptiloria exigua, 342
Puccinia Helianthi, 267
Pulsatilla, 490
Purshia, 331, 496, 497
Pustularia vesiculosa, 169
Pycnanthemum virginianum, 356, 363
Pyrenula, 191
Pyrethrum cinerarifolium, 126
Pyrola americana, 411, 415; chlorantha, 411, 415; elliptica, 29, 33, 34; rotundifolia, 29, 32, 34; uliginosa, 472
Pyrolaceae, 411
Pyronema, 158, 160, 161, 163, 164, 166, 167, 169, 171, 179, 180, 181, 184, 186, 188, 190, 192, 193, 194, 196; confluens, 159, 190
Pyrus coronaria, 362

Quamasia hyacinthina, 29, 35
Quercus alba, 213, 219; aquatica, 214; arkansana, 214; Catesbaei, 211, 559, 563, 566; cinerea, 211, 559, 563; falcata, 559; geminata, 559, 563; laurifolia, 213, 559; Margaretta, 211, 214; marylandica, 211, 214, 559, 563; Michauxii, 559; microcarya, 214; minor, 29, 34, 38; Morrisoniana, 298; myrtifolia, 214; nigra, 214, 219, 559; oblongifolia, 303; Phellos, 559; Prinus, 29, 33, 35; pumila, 563, 566; rubra, 219; severnensis, 298; sp. 214; stellata, 211, 559; velutina, 213; virginiana, 559
Quillaja Saponaria, 126

Radiovittaria, 391, 393, 397, 400
Radula arctica, 606, 607, 613, 615; Bolanderi, 606, 607, 609, 615; cochleariformis, 610; complanata, 613; Krausei, 613; **polyclada**, 607, 608, 609, 610, 615; prolifera, 609, 610; spicata, 607
Ramona incana, 341
Ranunculus, 466; adoneus, 101, 473; affinis, 96, 472; alismaefolius, 473; eximius, 473; glaberrimus, 339; hyperboreus, 97, 467; Macauleyi, 98, 468; nivalis, 98; pedatifidus, 96; pygmaeus, 93, 467; saxicola, 468; stenolobus, 101, 472, 473
Raritania gracilis, 298
Relationship between the number of ovules formed and the number of seeds developing in Cercis, On the, 243; Further observations, 533
Report on the Hepaticae of Alaska, 577
Review of investigations of the mosaic disease of tobacco, together with a bibliography of the more important contributions, A, 435
Revision of the genus Vittaria J. E. Smith, A, I. The species of the subgenus Radiovittaria, 391
Rhamnites apiculatus, 298
Rhexia Alifanus, 560, 565, 567; glabella, 567; lutea, 561; virginica, 74
Rhodiola, 466; integrifolia, 96, 99, 467; polygama, 99, 468; rosea, 96
Rhododendron maximum, 28, 29, 30, 32, 35, 41
Rhodotypus, 331
Rhus copallina, 560, 563; glabra, 363, 560, 563; glabra occidentalis, 341; mollis, 303; radicans, 214; Toxicodendron, 341, 363; Toxicodendron radicans, 366, 368
Ribes aureum, 340; montigenum, 465, parvulum, 465, 466
Riccardia, 599, 601; latifrons, 585; major, 612, 615; multifida, 586, 615
Riccia crystallina, 333; Frostii, 333; natans, 333

Ricciocarpus natans (L.) Corda. The appearance of polar bodies in the spermogenous tissue of, 333
Ricciocarpus natans, 333, 335
Ricinus, 230; communis, 107
Robinia, 548
ROBINSON, W. J. A taxonomic study of the Pteridophyta of the Hawaiian Islands—IV, 51
Roripa Nasturtium, 339
Rosaceae, Notes on, VII, 319; VIII, 483
Rosa humilis, 354, 357, 363; polyantha, 367, 370
Rubus idaeus aculeatissimus, 517; nigrobaccus, 560; occidentalis, 363, 367, 370; odoratus, 29, 34, 38; sp. 363
Rudbeckia hirta, 355, 366, 367, 512
Rumex Acetosella, 515, 516, 518, 519, 520 elongatus, 514, 515, 516, 518
RYDBERG, P. A. Notes on Rosaceae—VII, 319; VIII, 483; Phytogeographical notes on the Rocky Mountain region, II. Origin of the alpine flora, 89; III. Formations in the alpine zone, 459
Rydbergia Brandegei, 101; grandiflora, 101, 469, 470
Rynchospora axillaris, 561; Baldwinii, 561; Chapmani, 560; corniculata, 561; cymosa, 1; elongata, 2; glomerata, 30, 36; **jamaicensis**, 1; polyphylla, 2; semiplumosa, 561

Sabalites magothiensis, 298
Sabbatia angularis, 561; campanulata, 560; decandra, 567; gentianoides, 567; macrophylla, 560, 563, 567
Saccobolus, 193
Sagina nivalis, 97, 468; saginoides, 467
Sagittaria lancifolia, 561
Salix, 231, 472; alexensis, 102, 473; amygdaloides, 29, 35; anglorum, 99; arbusculoides, 102, 473; arctica, 99; Barrattiana, 102, 473; cascadensis, 99, 467, 468; chlorophylla, 95, 472; Dodgeana, 99, 467, 469; Drummondiana, 102, 473; flexuosa, 298; glaucophylla, 230; glauca, 95; glaucops, 95, 99, 472; Lesquereuxii, 296, 298; longifolia, 359; monticola, 472; nigra, 559; nivalis, 98, 467, 468; petrophila, 467; phyllicifolia, 95; pseudolapponum, 99, 468; reticulata, 93, 98; retusa, 99; saximontana, 98, 467, 472; Scouleriana, 338
Salsola Kali tenuifolia, 517
Sambucus canadensis, 362; glauca, 339; pubens, 29, 34; racemosa, 517
Sanguinaria, 548
Sanguisorba, 322, 323; canadensis, 322; latifolia, 322; media, 322; Menziesii, 322; microcephala, 322; officinalis, 322; sitchensis, 322
Sanicula canadensis, 83; marylandica, 83; sp. 214
Santalum album, 126
Sapotacites Knowltoni, 298
Saprolegnia mixta, 290
Sarcocolla, 223
Sarcopoterium, 323
Sarcoscyphus sphacelatus, 613
Sarracenia flava, 563, 566; psittacina, 561, 563, rubra, 566; Sledgei, 560, 563, 566
Sassafras acutilobum, 296, 297; varifolium, 560, 563
Satureja Acinos, 516
Sauteria alpina, 612
Saururus, 141, 229, 232; cernuus, 561
Saxifraga, 466; cernua, 94, 98, 467; debilis, 98, 468; pennsylvanica, 29, 35; rivularis, 98, 102, 468; simulata, 98; virginiensis, 29, 33, 35
Scapania albescens, 603; Bolanderi, 602, 615; cordifolia, 577, 603, 604, 605, 615; curta, 613; dentata, 603, 604, 605; imbricata, 613; irrigua, 606; nemorosa, 606, 615; paludosa 605; paludosa papillosa, 603, 605; planifolia, 616; uliginosa, 605, 606, 615; umbrosa, 606, 615; undulata, 603, 606
Schizonotus discolor, 339
Schoenocaulon officinale, 107, 127
Scilla, 224; maritima, 126
Scinaia, 177
Scirpus caespitosus, 472; cubensis, 1; lineatus, 359; pauciflorus, 472
Scleria Baldwinii, 561
Scleroderma, 179
Scutellaria galericulata, 29, 35; grandiflora, 341, 342; lateriflora, 107, 339
Sebastiana ligustrina, 214
Secale cereale, 516, 518, 520
Securinega, 230
Sedum acre, 514; stenopetalum, 339, 462, 463
Seed development in the genus Peperomia, 137, 221
Selaginella, 55, 57; arbuscula, 58, 59; deflexa, 57, 58; densa, 465, 466; flabellata, 58, 59; Menziesii, 58, 59; parvula, 59; Springii, 58, 59
Selaginellaceae, 57
Senecio, 466; amplectens, 471; atratus, 472; aurens, 97; Balsamitae, 358, 359; carthamoides, 465; crocatus, 468; cymbalarioides, 97, 467; Fremontii, 465, 466; Harbourii, 464; Holmii, 101, 468; hydrophilus, 339; invenustus, 465; lugens, 102; Nelsonii, 464; petrocallis, 101, 464, 466; pseudaureus, 470; Porteri, 101; Purshianus, 463; Soldanella, 101, 468; taraxacoides, 101, 468
Sequoia ambigua, 298; heterophylla, 298; Reichenbachi, 298

Serenoa, 563; serrulata, 559, 566
Sericocarpus asteroides, 29, 33, 35
Setaria viridis, 514, 515, 516
Sibbaldia procumbens, 321, 466, 467
Sidalcea candida, 470
Sieversia, 489, 490, 494; anemonoides, 489, 490, 491; brevifolia, 491, 493; calthifolia, 490, 493, 494; calthifolia congesta, 493; campanulata, 491; canescens, 491, 492, 493; ciliata, 339, 343, 370, 490, 491, 492, 493; flavula, 491, 493; glacialis, 491; grisea, 491, 492, 493; macrantha, 494; montana, 490, 493; Peckii, 92, 490, 493; pentapetala, 490, 491; radiata, 493; radicata, 490; rosea, 492; triflora, 491, 492, 493
Silene acaulis, 93, 463, 463; Douglasii multicaulis, 339; noctiflora, 514, 515, 517, 518; rotundifolia, 29, 30, 32, 35, 44, 45
Silphium, 358, 359; integrifolium, 355, 356; laciniatum, 354, 355, 363, 512; terebinthinaceum, 356, 357, 359, 363, 512
Siphonychia sp. 212
Sisymbrium altissimum, 514, 515, 518; officinale leiocarpum, 514, 515
Sisyrinchium sp., 357
Sium cicutaefolium, 84
Slope exposure as a factor in the distribution of Pseudotsuga taxifolia in arid parts of Washington, 337
SLOSSON, M. Notes on two North American ferns, 307
Smelowskia americana, 96, 462, 463; calycina, 96
Smilacina, 223, 231; stellata, 223, 224, 225
Smilax echirrata, 29, 36; laurifolia, 560; pumila, 214, 216
Solanum Dulcamara, 124, 339; nigrum, 514, 517, 519; tuberosum, 444
Solidago bicolor, 33; canadensis, 354, 357; ciliosa, 101, 470; decumbens, 101, 470; erecta, 30, 35; juncea, 29, 35; nemoralis, 354; odora, 212, 214; ohioensis, 359; oreophila, 101, 470; Riddellii, 359; rigida, 354, 356, 357; scopulorum, 101; serotina, 354
Sonchus alpinus, 624; asper, 517
Sorghastrum nutans, 354, 357, 360; secundum, 212
Sparganium angustifolium, 474; minimum, 474
Spartina Michauxiana, 358, 359; polystachya, 561
Spathularia, 180; stellaris, 98; Vreelandii, 98
Spermacoce floridana, 13; keyense, 13
Sphaeralcea rivularis, 340
Sphaerites raritanensis, 297
Sphaerosoma, 189
Sphaerotheca, 170, 182, 192, 193, 194; Castagnei, 159
Sphenolobus exsectus, 588; minutus, 588; saxicola, 613
Sphenopholis obtusata, 308
Spiraea corymbosa, 339
Spiranthes cernua, 356
Spirogyra, 267; communis, 290
Sporobolus heterolepis, 354, 356, 357, 360
Sporodinia grandis, 286
Stachys cordata, 29, 35; palustris, 339, 358
STANDLEY, P. C. New or notable species of Amaranthus, 505
Staphylea, 548
Stenophyllus ciliatifolius, 212
Stenostomum myrtifolium, 12
Steironema ciliatum, 339, 340
Sterculia cliffwoodensis, 298; minima, 298
Sterigmatocystis nigra, 290
Stigmatomyces Baeri, 161
Stipa spartea, 354, 356, 357
Stokesia laevis, 560, 563, 567
Strophanthus hispidus, 127; Kombe, 127; sp., 127
Strychnos Nuxvomica, 107, 126
Studies of West Indian plants—V, 1
Stylopus, 484; vernus, 484
Stylosanthes biflora, 29, 34, 38
Sullivantia Sullivantii, 29, 32, 35
Superficial study of the pine-barren vegetation of Mississippi, A, 551
Swertia palustris, 96, 472; perennis, 96
Symplocos tinctoria, 216, 219
Syntheris plantaginea, 465
Synthyris rubra, 339

Taeniopsis, 394
Taeniopteris, 394, 395
Taraxacum, 148, 149; arcticum, 98; officinale, 514, 515, 516, 517, 518, 519, 520, 521; scopulorum, 98, 466, 467
Taxodium ascendens, 565; distichum, 90, 559, 563, 565; imbricarium, 559, 563, 565
Taxonomic study of the Pteridophyta of the Hawaiian Islands—IV, A, 51
Tecoma radicans, 367, 369, 370; stans, 303
Teichospora, 196
Telesonix heucheriformis, 101, 465, 466; Jamesii, 101, 465
Tellima parviflora, 339; tenella, 339
Temnoma setiforme, 613
Tephrosia virginiana, 355
Thalictrum alpinum, 93, 470, 472; dasycarpum, 358, 359, 363
Thaspium aureum, 359
Thecotheus, 160, 190, 193; Pelletieri, 159, 192, 193
Thelebolus, 170, 179, 190, 194, 196; stercorarius, 159

Theobroma Cacao, 127
Thlaspi alpestre, 100; coloradense, 100, 462, 464; glaucum, 100; purpurascens 100, 462, 464
Thrinax microcarpa, 11
Thuidium sp. 214, 216
Thuja cretacea, 298; occidentalis, 512; plicata, 343
Thurberia thespesioides, 301, 302, 303, 304, 305
Tigarea, 496
Tilia heterophylla, 212, 219
Tillandsia usneoides, 212, 214, 560
Tium alpinum, 470
Tobacco, A review of investigations of the mosaic disease of, together with a bibliography of the more important contributions, 435
Tofieldia coccinea, 102; palustris, 95, 471; racemosa, 560
Tonestus Lyallii, 101; pygmaeus, 101, 468
Tradescantia reflexa, 356
Tremella, 179
Trichomanes hymenoides, 307; Krausii, 307; Petersii, 307; punctatum, 307
Trichostema dichotomum, 29, 35
Trifolium attenuatum, 101, 468; bracteolatum, 101, 468; Brandegei, 101, 467, 468; dasyphyllum, 101, 462, 463, 468; Haydeni, 471; hybridum, 514, 515, 516, 518, 519, 520; lividum, 101, 468; montanense, 101; nanum, 101, 470; Parryi, 101, 467, 470; pratense, 360, 514, 515, 516, 517, 518, 520; reflexum, 30; repens, 360, 514, 515, 516, 518, 519, 520; salictorum, 101; stenolobum, 101, 468
Trilisa odoratissima, 561
Trillium Hugeri, 214; petiolatum, 339
Triplasis americana, 212
Trisetum majus, 469; subspicatum, 93, 94, 463, 469
Trollius laxus, 30
Tsuga **americana,** 629; canadensis, 29, 33, 35, 45, 512, 621, 622, 629; heterophylla, 580; Mertensiana, 580
Tuber, 169, 179; excavatum, 182; puberulum, 182
TURESSON, G. Slope exposure as a factor in the distribution of Pseudotsuga taxifolia in arid parts of Washington, 337
Typha, 231; latifolia, 359

Umbelliferae, 83
Unifolium canadense, 29, 34, 36, 38
Usnea, The nomenclature of the genus, 373
Usnea, 169, 373; articulata, 374, 376, 377; barbata, 159, 373, 374, 376, 377; barbata articulata, 373; barbata cera; tina, 373; barbata dasypoga, 373a barbata florida, 373; barbata florida hirta, 373; barbata florida rubiginea, 373; barbata florida strigosa, 373; barbata hirta, 373; barbata intermedia, 378; barbata plicata, 373; barbata rubiginea, 373; barbata **stricta,** 378; californica, 376; ceratina, 374, 375, 376, 377, 378; coralloides, 376; dasypoga, 374, 376; dasypoga plicata, 374, 377; florida, 374, 375, 377; florida hirta, 375; plicata, 374, 375, 377; plicata barbata, 374; plicata dasypoga, 377
Utricularia vulgaris, 474

Vacciniaceae, 419
Vaccinium, 423; amoenum, 423; angustifolium, 420, 421, 422, 423, 424; angustifolium × corymbosum, 424; **atlanticum,** 422, 423; atrococcum, 425, 426; australe, 424; **Brittonii,** 420, 421, 422; caesariense, 424; caespitosum, 97, 467; canadense, 421; corymbosum, 420, 423, 424, 425, 426; corymbosum amoenum, 423; corymbosum glabrum, 424; disomorphum, 425; Dobbini, 422; fuscatum, 426; Myrtillus, 96; nigrum, 420; oreophilum, 96, 470, 472; pennsylvanicum, 420, 421; pennsylvanicum nigrum, 420, 421; scoparium, 470; vacillans, 420, 422; **vicinum,** 425; virgatum, 423
Vagnera amplexicaulis, 339; stellata, 415
Valeriana acutiloba, 470; edulis, 359; officinalis, 127; pauciflora, 29, 35; septentrionalis, 471
Valerianella macrocera, 339
Varronia **clarendonensis,** 16
Vaucheria, 167
Vauquelinia californica, 303
Veratrum album, 126; nigrum, 126; viride, 126; Woodii, 30
Verbascum Thapsus, 514, 515, 516, 517, 518, 519, 520, 521
Verbena bracteosa, 512; stricta, 354
Verbesina, 362
Veronica virginica, 358, 363; Wormskjoldi, 472
VESTAL, A. G. Black-soil prairie station in northeastern Illinois, A, 351
Viburnum dentatum, 29; molle, 29; nudum, 560; rufidulum, 214
Vicia americana, 339
Viola adunca, 339; biflora, 90; hirsutula, 29, 32, 35; papilionacea, 358; pedatifida, 356, 357; retroscabra, 339; rostrata, 29, 35; rotundifolia, 29, 32, 35; sp. 214
Vitis aestivalis, 560; rotundifolia, 214; vinifera, 367, 369; vulpina, 362, 363
Vittaria J. E. Smith, A revision of the genus, I. The species of the subgenus Radiovittaria, 391
Vittaria, 391, 392, 393, 394, 395, 399, 400, 404, 406; Bommeri, 401, 407;

elongata, 394; Forbsei, 394; Gardneriana, 398, 400, 401, 404, 407; gracilis, 400, 401; Karsteniana, 400, 401; **latifolia**, 398, 403, 404, 407; lineata, 394, 395, 397, 404; longipes, 405, 406; minima, 398, 399, 400; Moritziana, 405, 406; Orbignyana, 405, 406; remota, 398, 401, 402, 403, 404; Ruiziana, 398, 405; scolopendrina, 394, 395; sikkimensis, 395, 399; stipitata, 398, 404, 406; **Williamsii**, 398, 407

Waldsteinia, 331; Doniana, 331, 332; Fragarioides, 331, 332; idahoensis, 332; lobata, 332; parviflora, 332
Warea cuneifolia, 212
West Indian plants, Studies of,—V., 1
Widdringtonites Reichii, 298
Wild cotton plant (Thurberia thespesioides) in Arizona, The, 301
Williamsonia delawarensis, 297; marylandica, 297
Wisteria frutescens, 367, 368
Wrangelia, 173, 178

Xolisma ligustrina, 414
Xylaria, 170; tentaculata, 159
Xylorrhiza Brandegei, 101, 464; coloradensis, 101, 464
Xyris flexuosa, 561; sp. 561

Zodiomyces vorticellarius, 161
Zygalchemilla, 319, 321
Zygia, 8; arborescens, 8